

VDAC1-5 | Voltage-dependent anion-selective channel protein 1-5, Biotin-conjugated (40 µg)

Cat PA01192

Size 40 µg

Host

Rabbit

Clonality

Polyclonal

Confirmed reactivity

Arabidopsis thaliana, Beta vulgaris, Brassica oleracea var. botrytis, Oryza sativa, Papaver sp. pollen tubes (IL), Spinacia oleracea, Physcomitrium patens

Immunogen

KLH-conjugated peptide conserved in all known higher plant VDAC proteins including Arabidopsis thaliana VDAC1 UniProt: Q9SRH5, TAIR: AT3G01280, VDAC2 UniProt F4K3R8-1, TAIR: AT5G67500, VDAC3 UniProt: Q9SMX3-1, TAIR: AT5G15090, VDAC4 UniProt: Q9FKM2-1, TAIR: AT5G57490, VDAC5 UniProt: Q9M2W6-1, TAIR: AT3G49920

Host

Rabbit

Clonality

Polyclonal

Purity

Immunogen affinity purified serum in PBS pH 7.4, conjugated to biotin.

Format

Liquid

Storage

Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years.

Application

2D Blue Native PAGE (2D BN-PAGE), Immunolocalization (IL), Western blot (WB)

Recommended dilution

1 : 500 (IL), 1 : 2000, 2-30 µg protein/lane (WB)

Expected | apparent MW

29 kDa (for Arabidopsis thaliana)

Confirmed reactivity

Arabidopsis thaliana, Beta vulgaris, Brassica oleracea var. botrytis, Oryza sativa, Papaver sp. pollen tubes (IL), Spinacia oleracea, Physcomitrium patens

Predicted reactivity

Arabidopsis alpina, Aundo donax, Brachypodium distachyon, Brassica campestris, Brassica napus, Brassica rapa subsp. pekinensis, Capsella rubella, Citrus clementina, Citrus sinensis, Eutrema salsugineum, Glycine max, Glycine soja, Gossypium arboreum, Hoedum vulgare var. distichum, Jatropha curcas, Medicago truncatula, Mesembryanthemum crystallinum, Morus notabilis, Nicotiana tabacum, Phaseolus coccineus, Phaseolus

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY

VDAC1-5 | Voltage-dependent anion-selective channel protein 1-5, Biotin-conjugated (40 µg)

Cat PA01192

Size 40 µg

vulgaris, Pisum sativum, Plantago major, Prunus persica, Ricinus communis, Solanum lycopersicum, Solanum tuberosum, Sorghum bicolor, Theobroma cacao, Triticum aestivum, Vitis vinifera, Zea mays

Not reactive in

Chlamydomonas reinhardtii, Glycine max, Zea mays, diatoms, Saccharomyces cerevisiae

Additional information

Cellular [compartment marker] of mitochondrial outer membrane, Amount of mitochondrial fraction detected by anti-VDAC1 antibody is from 2-10 µg. Immunolocalization method description and images are available here Blue-native (2D BN/SDS-PAGE) methodology is described in Piechota et al. 2010

Description

VDAC proteins are porin-type, β -barrel diffused pores. It is mainly located in the outer membrane of mitochondria and participates in metabolite exchange.

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY