

Cytokinin | Dihydrozeatin riboside (for immunolocalization)

Cat PA01619

Size 200 µg

Host

Rabbit

Clonality

Polyclonal

Confirmed reactivity

Dihydrozeatin riboside

Immunogen

BSA-conjugated, via ribose, dihydrozeatin riboside

Host

Rabbit

Clonality

Polyclonal

Purity

Total IgG. Protein G purified in PBS.

Format

Liquid

Storage

Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application

Immunolocalization (IL)

Recommended dilution

Specific information about dilution is going to be included on the vial

Confirmed reactivity

Dihydrozeatin riboside

Predicted reactivity

Dihydrozeatin riboside

Not reactive in

No confirmed exceptions from predicted reactivity are currently known

Additional information

Cytokinin | Dihydrozeatin riboside (for immunolocalization), For 1 mg of antibodies For ELISA and immunoassay or For ELISA kit - please inquire, Antibodies are supplied in 50% glycerol

Description

Cytokinins (CK) are a class of plant growth substances (plant hormones) that promote cell division through local

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY

Cytokinin | Dihydrozeatin riboside (for immunolocalization)



Cat PA01619

Size 200 μ g

and long-distance signaling. Zeatin is an adenine-type cytokinin synthesized in stems, leaves and roots.

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY
