# Phaseolus vulgaris Lectin (PHA-E+L) - Ferritin



Cat PL00598

Size 1 mg

## **Sugar Specificity**

Complex N-Glycans

# **Description**

Phaseolus vulgaris lectin (phae +L) is coupled to ferritin and has a molecular weight of 474,000 g/mol(all 24 subunits combined). Ferritin binding lectins can be used to visualize the location of specific carbohydrate parts and to examine the distribution of plasma membrane glycoproteins. These conjugates can be observed by electron microscopy.

## **Abbreviation**

PHA-E+L

### **Material Source**

Kidney bean

## Conjugate

Ferritin

#### Concentration

1 mg/mL

## **Purity**

High Purity Grade

#### **Shelf Life**

1 year

# **Blood Group Specificity**

Non-specific

# **Sugar Specificity**

Complex N-Glycans

# **Inhibiting or Eluting Sugar**

Bovine Thyroglobulin, Acetic Acid

#### **Divalent Ions**

Ca++, Mn++

# **Mitogenic Activity**

Yes

### **Format**

Liquid

## **Storage**

-20°C

# **Hazardous Shipping**

Non-hazardous

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY

Address: SUITE 209, 17 Ramsey Road, Shirley, NY 11967 Tel: 1-631-637-0420

# Phaseolus vulgaris Lectin (PHA-E+L) - Ferritin



Cat PL00598

Size 1 mg

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY

Address: SUITE 209, 17 Ramsey Road, Shirley, NY 11967 Tel: 1-631-637-0420