

GST conjugated to DyLight 650

Cat PA00165

Size 100 µg

Host

Goat

Clonality

Polyclonal

Confirmed reactivity

Immunogen

Glutathione-S-transferase from *S. japonicum*

Host

Goat

Clonality

Polyclonal

Purity

Immunogen affinity purified using solid phase GST in 100 mM NaPO₄ (pH 7.4), 100 mM NaCl, 2 mM sodium azide.

Format

Lyophilized

Reconstitution

For reconstitution add 100 µl of sterile water

Storage

Store at 2-8°C.; Product is stable for up to 4 weeks at 2-8°C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and Store at -20°C.

Application

Flow Cytometry (FL), Cell Based Assays (CBA)(CBA)

Recommended dilution

The optimal working dilution should be determined by the investigator

Additional information

Antibody is provided in: 100 mM Sodium Phosphate (pH 7.4), 100 mM Sodium Chloride, 2 mM Sodium Azide

Description

Goat anti-GST IgG is a primary antibody that binds to GST and is directly coupled to DyLight 650. This is due to the advantage of shortening the assay time by not requiring the use of GST secondary antibodies. DyLight is a trademark of Thermo Fisher Technologies Co., Ltd. and its subsidiaries.

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY